

2017-03-31

Advances in physiological and molecular aspects of plant cold tolerance

Rihan, Hail

<http://hdl.handle.net/10026.1/10527>

10.1080/17429145.2017.1308568

Journal of Plant Interactions

Informa UK Limited

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.



Advances in physiological and molecular aspects of plant cold tolerance

Hail Z. Rihan, Mohammed Al-Issawi & Michael P. Fuller

To cite this article: Hail Z. Rihan, Mohammed Al-Issawi & Michael P. Fuller (2017) Advances in physiological and molecular aspects of plant cold tolerance, Journal of Plant Interactions, 12:1, 143-157, DOI: [10.1080/17429145.2017.1308568](https://doi.org/10.1080/17429145.2017.1308568)

To link to this article: <http://dx.doi.org/10.1080/17429145.2017.1308568>



© 2017 The Author(s). Published by Informa UK Limited, trading as Taylor & Francis Group



Published online: 31 Mar 2017.



Submit your article to this journal [↗](#)



View related articles [↗](#)



View Crossmark data [↗](#)

REVIEW ARTICLE



Advances in physiological and molecular aspects of plant cold tolerance

Hail Z. Rihan^a, Mohammed Al-Issawi^b and Michael P. Fuller^c

^aFaculty of Agriculture, University of Damascus, Damascus, Syria; ^bAgriculture College, University of Anbar, Ramadi, Iraq; ^cSchool of Biological & Marine Sciences, Plymouth University, Plymouth, UK

ABSTRACT

Abiotic stress is one of the main causes of crop reduction globally. Among the different abiotic stresses, cold is an essential factor that limits crop productivity worldwide. Low temperature affects the growth, development and distribution of agronomic species throughout the world. Cold stress is a serious threat to the sustainability of crop yields. Indeed, cold stress can cause major crop losses. A significant number of researches have been reviewed and discussed in this study in order to improve the understanding of the physiological and genetic nature and function of plant cold tolerance. Recent developments in determining the mechanism of genes with roles in freezing tolerance and the systems involved in low-temperature gene regulation and signal transduction are described. The roles of a family of *Arabidopsis* transcription factors, the *CBF/DREB1* proteins, have been described and its role in controlling the expression of a regulon of cold-induced genes (COR) that increase plant freezing tolerance has been explained. Moreover, this study has reviewed the recent application applied to improve the cold tolerance of plants such as molybdenum. The use of infrared camera to study the process of plant injuries caused by low temperature has also been reviewed.

ARTICLE HISTORY

Received 5 January 2017
Accepted 15 March 2017

KEYWORDS

Cold tolerance; CBF/DREB;
COR genes; infrared
thermography; ABA

Plant abiotic stress

Plant growth and productivity are limited by both biotic and abiotic factors (Seki et al. 2003). A series of morphological, physiological, biochemical and molecular changes can be triggered in plants under the effect of abiotic stresses which eventually adversely affect the growth and productivity of crops. Therefore, plants respond and adapt in order to survive at the molecular and cellular levels in addition to physiological and biochemical responses (Sanghera et al. 2011). It was reported that abiotic stresses can cause a decrease in yield of crops up to 50% resulting in very high economic losses (Vij & Tyagi 2007). The world population is increasing rapidly and at the same time food production is decreasing due to the effect of various abiotic stresses (Mahajan & Tuteja 2005). Therefore, the minimization of these losses is one of the main objectives for plant and crop specialists and since it is very difficult to control abiotic stress resulting from climate change and human activities, the development of stress-tolerant crop genotypes is necessary (Mahajan & Tuteja 2005).

Different scientists have defined stresses differently depending on their mode of studies. Lichtenthaler (2006) defined stress as any substance or condition which stops a plant's metabolism, development or growth. Stress was also defined by Mahmood (2002) as any element that decreases plant reproduction and development below the genotype's potential. However, plant response to stress is very complex, including morphological, physiological, biochemical and molecular changes. Some plants have morphological adaptations in order to avoid undesirable conditions and others can alter their physiology metabolism, gene expression and various developmental activities to tolerate the effect of specific stresses.

Low-temperature stress (frost damage)

Among the different abiotic stresses, cold is an essential factor that limits crop productivity worldwide. Low temperature affects the growth and development of agronomic species throughout the world (Pearce 2001). The survival of plants in freezing temperatures depends on their capacity for cold acclimation (McKhann et al. 2008). Low temperature limits the geographical distribution of plant species and significantly decreases the yield of several crops around the world (Pearce & Fuller 2001). It is very important to study the frost damage mechanism and to breed cold-tolerant varieties since the average minimum temperature is below 0°C in about 64% of the earth's land area and it is below -10°C in about 48% (Deane 1994).

Frost technically means the formation of ice crystals and this includes a phase change from vapor or liquid to solid. Ice formation (crystallization) consists of two main steps: (a) nucleation; (b) crystal growth. Nucleation is the process when the molecules start to gather, forming clusters at the nanometer scale. However, these clusters are required to grow to a specific size in order to form stable nuclei and at this point the atoms arrange in a defined manner to form the crystal structure. Subsequently, the nuclei keep growing to form ice and cause damage effects on plant cells (Sakai & Larcher 1987).

Research about the mechanism of ice initiation in plants has greatly decreased recently because of its essential role in specific aspects of cold hardiness (Gusta et al. 2009; Aryal & Neuner 2010; Hacker et al. 2011; Walters Jr et al. 2011). Ice formation may take place either in the intracellular or extracellular spaces depending on the cooling conditions (Guy 1990). Intercellular freezing can happen when cooling

rates are high or after a significant supercooling. Supercooling is defined as when liquid water cools below the melting temperature (0°C) without any ice formation. The supercooling point is the minimum low temperature reached before ice formation. Extracellular freezing takes place in the spaces between cells in water transporting elements (xylem and phloem) or on the external surfaces of the plant. Ice will grow and spread from the initial formation point (nucleation site) through the extracellular spaces in contiguous water. Moreover, as long as the plasma membrane is unharmed and the cooling rate is low, the ice will stay in the extracellular spaces and it will draw the water from the cells since water has a lower osmotic potential in the ice (solid) phase than that in the liquid phase. This process will continue until equilibrium of water potential is achieved resulting in a dehydration effects on the plant cells (Wisniewski & Fuller 1999). Several forms of membrane damage can result as a consequence of dehydration caused by freezing effects. This could include expression-induced lysis, lamellar-to-hexagonal-phase transitions and additional jump lesions (Steponkus et al. 1993). McKersie and Bowley (1997) reported that freezing contributes to membrane damage through the induction of reactive oxygen species (ROS) production. It has been demonstrated that protein denaturation caused by the effect of low temperature could cause cellular damage (Guy et al. 1998). Takahashi et al. (2013) reported that the alternation of plasma membrane compositions and functions during cold acclimation is one of the most important adaptation mechanisms for plant cold tolerance.

Generally, plants can be classified into three different classes according to their low-temperature tolerance (Stushnoff et al. 1984). The first group includes frost-tender plants that are sensitive to chilling injury and can be killed by short periods of exposure to temperature just below freezing point. Beans (*Phaseolus lunatus*), corn (*Zea mize*), rice (*Oryza sativa*) and tomatoes (*Solanum lycopersicum*) are examples of plants in this category. Low-temperature acclimation of plants in the second group allows them to tolerate the presence of extracellular ice in their tissues. Their frost resistance ranges from the broad-leaved summer annuals, which are killed at temperatures slightly below freezing point, to perennial grasses that can survive exposure to -40°C. As temperatures decrease the outward migration of intracellular water to the growing extracellular ice crystal causes dehydration stress that will eventually result in irreversible damage to the plasma membrane. The final group is made up of very cold hardy plants that are predominantly temperate woody species. Like the plants in the previous group, their lower limits of cold tolerance are dependent on the stage of acclimation, the rate and the degree of the temperature decline and the genetic capability of tissues to accommodate extracellular freezing and the accompanying dehydration stress. Deep supercooling allows certain tissues from plants in this group to survive low temperatures without formation of extracellular ice. However, plants have adapted two mechanisms to protect themselves from damage due to below freezing temperatures. First, supercooling is a low-temperature tolerance mechanism that is usually associated with acclimated xylem parenchyma cells of moderately hardy woody plants. The second and most common low-temperature response mechanism is acclimation. Acclimation is a gradual process during which there are changes in just about every measurable morphological, physiological and biochemical characteristic of

the plant. These changes are determined by genotype and environmental interactions that are quite complex (George et al. 1982).

Cold acclimation and frost stress tolerance

Cold acclimation is defined as the exposure of plants to low non-freezing temperatures leading to significant positive effect on the cold tolerance of the plants (Thomashow 2010; Pearce et al. 2013). Extensive researches have been conducted to improve the understanding of the biochemical and molecular basis of the cold acclimation response and the changes that take place throughout this process (Lissarre et al. 2010). However, the increase in cold tolerance obtained by acclimation is not static. It depends on the season and it is lost quickly (deacclimation) when plants are exposed to warm temperatures. Wanner and Junttila (1999) also reported that cold acclimation is a photosynthesis-activity-demanding process. The authors reported that light plays a role in cold acclimation and moderate to high light intensity is required for a sufficient cold acclimation process. Ivanov et al. (2012) reported that the strength of light and the ambient temperature could be integrated into electron sink alternation in chloroplasts inducing a signal transduction pathway. They reported that the energy balance in chloroplasts could play a role in sensing ambient temperature. However, a plastid signal contributes to the rhythm of CBF expression and downstream of cold-responsive genes expression during the day/night cycle (Norén et al. 2016). Moreover, cold-tolerant species such as winter wheat, spinach, etc. have the capacity to maintain a high level of CO₂ assimilation rate whereas sensitive cold species do not have this capacity (Yamori et al. 2010). Gusta and Wisniewski (2013) reported that energy is an essential factor to drive acclimation and that sugars have important effects on plant freezing tolerance, resiliency of photosynthesis under stress condition. Dahal et al. (2012) reported a significant correlation between the accumulation of sugars and the development of cold tolerance in plants. The timing of cold acclimation and deacclimation, the onset and release from dormancy, as well as the timing of bud break play an integral role in the life cycle of woody plants and their adaptation to the external environment (Ríos et al. 2014; Wisniewski et al. 2015).

Extensive physiological and biological changes occur during cold acclimation (Figure 1) starting with a reduction in the growth rate and water content of various plant tissues (Levitt 1980). Through the cold acclimation process, reprogramming of gene expression and various modifications in the metabolism take place (Chinnusamy et al. 2010). Acclimation also causes an increase in the production of antioxidants, abscisic acid (ABA) and compatible osmolytes such as soluble sugars and proline (Lynch & Steponkus 1987; Koster & Lynch 1992; Chen et al. 1993; Kishitani et al. 1994; Uemura & Steponkus 1994; Murelli et al. 1995; Nomura et al. 1995; Dörffling et al. 1997; Tao et al. 1998). Denesik (2007) reported that cold treatment affects membrane fluidity resulting in an increase in the membrane rigidity. This more fluid state of cell membranes helps to protect cells under the effect of low-temperature stress through maintaining the cell shape and preventing cellular component from water loss. Moreover, the rigidity of membrane could reduce cells collapsing during extracellular freezing by creating a negative pressure in the cells (Rajashekar & Lafta 1996; Heidarvand




Figure 1. Cold acclimation induces changes in cellular processes. Different responses are observed while exposing plants to low non-freezing temperatures. Modified from Xin and Browse (2000).

& Maali Amiri 2010). Takahashi et al. (2013) reported that the alternation of plasma membrane composition and function is one of the most important mechanisms of plant cold tolerance. A proteome profiling to characterize alternations of the *Arabidopsis* microdomain during cold acclimation has been conducted by Minami et al. (2009) who reported the increase of clathrins and dynamin-related proteins in the microdomain during cold acclimation. Many proteins such as P-type ATPases, aquaporins and tubulins have been reported to accumulate under the effect of low-temperature treatment (Minami et al. 2009; Takahashi et al. 2013) (Figure 2). For more details, please read Takahashi et al. (2013).

Cold acclimation leads to large-scale transcriptome changes that program the production of an array of proteins to help plants surviving freezing stress (Kocsy et al. 2010; Winfield et al. 2010). Recent studies describing full genome transcripts and mutational and transgenic plant analysis have provided a great deal of information about the complex transcriptional system that functions under cold acclimation (Jan et al. 2009). Various regulatory pathways were found to be involved in the cold response when transcriptome profiling of around 8000 genes of *Arabidopsis* was carried out. It was observed that 300 genes were affected by cold, of which 218 showed an increase of transcripts expression and 88 showed a decrease of transcripts expression within 7 days of cold treatment (Fowler & Thomashow 2002). The use of microarray systems has enabled researchers to demonstrate that a large number of genes are induced under abiotic stresses (Seki et al. 2001; Kreps et al. 2002; Bray 2004; Maruyama et al. 2004; Vogel et al. 2005; Jan et al. 2009). The genes affected by low temperature have been classified into two main groups (Seki et al. 2002): the first group involves the proteins that function as a response to cold stress such as LEA proteins (Cushman & Bohnert 2000); the second group contains proteins which have roles in the further regulation of gene expression in cold stress conditions. Some of these pathways are considered to have multiple functions since they have also been reported to be involved in stress responses to drought and high salinity (Seki et al. 2003).

The majority of studies to improve the cold tolerance of plants have been conducted with the *Arabidopsis* plant (Gery et al. 2011). However, Li et al. (2012) reported that *Brachypodium distachyon* could be an important model to

investigate the molecular mechanism of low-temperature tolerance in core Pooideae species since *B. distachyon* have cold-responsive IRIP. Moreover, a large C repeating binding factor genes CBF3 subfamily was identified in *B. distachyon*. The authors also confirmed that FST-motifs and cold treatment that increased fructan accumulation were intensely different in *B. distachyon* compared to core Pooideae species.

Gene expression induction in response to low temperature

Cold acclimation affects the expression of a huge number of genes by either up- or down-regulations (Hannah et al. 2005). Furthermore, there is crosstalk between the response of different abiotic stresses and some of the genes up-regulated by drought or salt stress (Thomashow 1999; Seki et al. 2003; Zhao & Zhu 2016). Fowler and Thomashow (2002) reported that cold acclimation activates different gene expression pathways which in turn means that cold-induced genes could be members of more than one cold regulon. However, it should be mentioned that plant response to abiotic stresses is mediated by a common group of reactions, cooperatively known as signal transduction (Heidarvand & Maali Amiri 2010). Therefore, plant response to low temperature goes through specific process starting by cell recognition of low temperature (sensing) followed by the signaling process in order to induce the cold-responsive genes resulting in an increase in plant cold tolerance (Denesik 2007).

Free Ca^{2+} has been observed to increase rapidly in the cytoplasm of plant cells during the acclimation process and, therefore, it is most likely to be involved in this process (Eckardt 2009). Protein phosphorylation seems to be involved since the treatment of alfalfa plants with the protein phosphatase inhibitor okadaic acid increased the transcript levels of the alfalfa cold-responsive cas15 gene at normal temperature and the transcript level of this gene did not increase to normal levels when plants were exposed to low-temperature treatment after they were treated with the protein kinase inhibitor staurosporine. Furthermore, cold effects significantly decreased the protein phosphatase 2A activity which has been reported to be dependent on calcium influx. Thus, calcium is believed to prevent phosphatase 2A activity leading to the phosphorylation of one or more




Figure 2. The adaptation mechanism of the plasma membrane during cold acclimation reported by proteomics studies (Takahashi et al. 2013).

proteins that could have parts in the induction of genes involved in cold tolerance and acclimation (Monroy et al. 1998 cited from Thomashow 2001).

Knight and Knight (2000) indicated that the free cytoplasm Ca^{2+} observed under low-temperature effects comes from extracellular and intracellular calcium stores. Significant evidence have confirmed the essential role of Ca^{2+} signaling on cold induction of CBF pathways (Knight & Knight 2000). Ca^{2+} is considered to be the main signal transducer in signaling cascades motivated in response to plant abiotic stress types. Ca^{2+} also seems to be the crosstalk between abiotic stress pathways since the level of this cation increases under the effects of different types of abiotic stresses. Ca^{2+} binding proteins (Ca^{2+} sensor) mediate the signal activated by Ca^{2+} influx under the effect of abiotic stresses. Jenks and Wood (2010) reported three main Ca^{2+} proteins in plants: Ca^{2+} -dependent proteins kinases, calmodulin (CaM) and calcineurin B-like proteins. Rudd and Franklin-Tong (2001) reported that CaM, which a highly conserved protein, is the first sensor to the Ca^{2+} cytosolic level in all eukaryotic cells. Therefore, calmodulin genes, which control calcium fluxes, are considered to be the main sensors of environmental triggers such as extreme temperatures. Calmodulin is a calcium-binding protein which binds to a regulatory element in the *CBF2* gene promoter which plays a role in controlling the CBF regulon and freezing tolerance (Doherty et al. 2009). Eckardt (2009) reported that CAMTA play a significant part in transduction of low-temperature-induced cytosolic Ca^{2+} signals into downstream regulation of gene expression. However, Marc et al. (2010) reported clear evidence that physical changes in membrane fluidity initiated by the stoichiometry of lipids could be the main trigger for calcium influx into the cell.

Nuclear Ca^{2+} is increased after cold treatment and the maximum increase is delayed at 5–10 s compared to the peak of cytosolic calcium (Miura & Furumoto 2013). Mauger (2012) reported that the increase in nuclear Ca^{2+} is caused by a nuclear envelope that is considered one of the major Ca^{2+} stores. Moreover, they indicated that the increase in nuclear

Ca^{2+} can be amplified by cytosolic Ca^{2+} transients through the nuclear pore complex. Because of the similarity in pore complex system in both animal and plant nuclear membrane (Xu & Meier 2008), it was suggested that the nuclear Ca^{2+} signal could be started from the nuclear envelope and propagated by cytosolic Ca^{2+} transients in plants as well. However, nuclear Ca^{2+} has been reported to have an important role in controlling gene transcription in both plants and animals (Mazars et al. 2011; Mauger 2012).

ROS also play an important role as second messengers responding to various abiotic stresses (Tyystjärvi 2013). Rao et al. (2006) reported that abiotic stresses cause an oxidative burst and that a low level of ROS induces an increase in Ca^{2+} influx into the cytoplasm. The high level of Ca^{2+} activates NADPH oxidase in order to produce ROS through yielding O_2^- which is then converted to H_2O_2 (ROS) under the effect of super oxidase dismutase (SOD). Therefore, the production of ROS is Ca^{2+} dependent and the concentration of Ca^{2+} is also regulated by the concentration of ROS by the activation of Ca^{2+} channels in the plasma membrane (Kwak et al. 2003). Therefore, a crosstalk between Ca^{2+} and ROS modulates the activity of specific proteins that control the expression-specific definitive defense genes in the nucleus. However, several other genes have been demonstrated to have roles in the abiotic stress responses such as kinases and transcription factors which in turn have roles in the crosstalk between signaling cascades involved in responses against two or more kind of stresses (Rao et al. 2006).

Kocsy et al. (2008) reported a crucial role of glutathione in the ascorbate-glutathione cycle. They reported that glutathione has an important role in the regulation of the concentrations of hydrogen peroxide (H_2O_2) in plants. When plants are exposed to low temperature (0–15°C), chilling and cold acclimation, an increase in H_2O_2 level is observed. The accumulation of H_2O_2 is a result of the reduction of oxygen by the excess electrons coming from the photosynthetic and respiratory electron transport chains under low-temperature conditions. This leads to the accumulation of superoxide radicals (O_2^-) which are subsequently converted to form H_2O_2

via SOD. H_2O_2 has a damaging effect on the cell membrane and protein structures of the cells. Therefore, plants developed an effective mechanism for the removal of excess H_2O_2 through the conversion of reduced glutathione (GSH) to its oxidized form (GSSG). However, GSH is reproduced under the effect of NADPH-dependent glutathione reductase (GR). The high level of GSH and GR observed under the effect of low temperature has been considered as an important indicator of a possible contribution of these materials to low-temperature tolerance in plants (Kocsy et al. 2008). It has been reported that the GSH/GSSG ratio and changes in H_2O_2 level can have an important role in the cold acclimation resulting in an activation of redox state of the cells and this in turn could activate special defense machines via a redox signaling chain under the effect of acclimation or low temperature. This hypothesis has been supported by the discovery of several defense genes that have antioxidant-responsive elements or GSSG-binding sites in their regulatory regions (Kocsy et al. 2008).

Ghildiyal and Zamore (2009) indicated that micro-RNAs (miRNAs) and small interfering RNAs could be repressors of gene induction in both plants and animals. Many cold-regulated miRNAs have been recognized using bioinformatics, cloning and sequencing tools in different plant species (Zhou et al. 2008; Zhang et al. 2009; Lv et al. 2010; Chen et al. 2012). However, only a little is known about the mi target genes that are unregulated under the effect of low temperature (Miura & Furumoto 2013).

Although a large number of studies have investigated the cold-tolerance mechanism in plants, there is still a lack in the progress of improving the cold frost tolerance of crop plants and this could be caused by difficulty of synthetically creating the complex environmental conditions existing when plants acclimate in nature (Gusta et al. 2009). Plants acclimated in growth chambers might respond differently from those growing in nature since the growth conditions such as light, air and soil temperature and temperature variations are different between the growth chamber and the field (Gusta & Wisniewski 2013).

CBF cold-response pathway

CBF gene discoveries are the most significant in the field of low-temperature adaptation and signal transduction. CBFs are discovered in all important field crops and some vegetable species (Sanghera et al. 2011). CBF genes have been identified in a wide range of herbaceous (Mizoi et al. 2012) and woody plants (Wisniewski et al. 2014). Intensive studies in the expression of cold-response gene expression in *Arabidopsis* resulted in the identification of the *CBF/DREB1* transcription factors family which have an essential role in cold acclimation and freezing stress tolerance in plants (Gilmour et al. 1998). These transcription factors bind to specific regulatory sequences in the promoters of cold and dehydration-responsive genes. These sequences are C-repeat (CRT: TGGCCCG AC) and dehydration-responsive elements (DRE: TACCG ACAT). Both of these sequences contain the highly conserved core 5-bp sequence of CCGAC, which has the capacity to regulate transcription under drought, low temperature and salinity (Baker et al. 1994; Yamaguchi-Shinozaki & Shinozaki 1994; Gao et al. 2007). Therefore, *CBF* induces the expression of *COR* genes (the genes which contain the *COR* sequence) and these genes play an essential role in the improvement

of plant cold tolerance (Mizoi et al. 2012). *CBF/DREB1* is a family of transcription factors that belongs to the ERF/AP2 transcription family (Yamaguchi-Shinozaki & Shinozaki 2005). The finding of Yamasaki and Randall (2016) has confirmed the initial steps in the *CBF/DREB1* pathway in soybean but the contributions to the lack of *GmCBF/DREB1* transcripts and/or protein, which are not sufficient to up-regulate downstream genes, absence or co-activators or presence of strong negatively acting transcriptional regulators or lack of appropriate *CBF/DREB1*-responsive element in the promoters of critical cognate soybean genes. Thomashow (2001) has characterized three different cold-inducible *CBF/DREB1* genes in *Arabidopsis* described as: *CBF/DREB1B*, *CBF2/DREB1C* and *CBF3/DREB1A*. *CBF3/DREB1A* and *CBF/DREB1B* are induced at the same time and earlier than *CBF2/DREB1C* (Medina et al. 2011). More recently Monroe et al. (2016) screened 477 wild accessions from *Arabidopsis* for the variation in CBF genes; they found that CBF sequences variation is strongly associated with winter temperatures. The types that come from warmer winters showed significant access to the CBF polymorphisms. *RNAi* lines targeting all three *CBF* genes in *Arabidopsis thaliana* were obtained in eight different accessions in order to evaluate the quantitative influence of *CBF* expression on plant freezing tolerance. Outstanding differences were observed between different accessions in terms of the effects that reduced CBF expression had on freezing tolerance, though effects on growth were mostly very minor to draw steady conclusions (Gery et al. 2011). The authors reported that the analysis of the relationship between different types of CBFs revealed a tight co-regulation between CBF1 and CBF3, whereas the relationship between CBF2 and CBF1 or CBF3 expression levels was intensely related to the genetic background of the *RNAi* lines. However, while three closely related *CBF* genes have been reported in *Arabidopsis*, wheat and some other cereals have undertaken an extension in this family (Pearce et al. 2013). Badawi et al. (2007) reported that there are at least 15 *CBF* genes in each of the three hexaploid wheat genomes.

The *CBF/DREB1* over-expression transcriptome analysis in transgenic plants demonstrated that about 12% of *COR* genes are controlled by *CBF/DREB1* in *Arabidopsis*. However, among the three CBF factors no specific target was observed (Zeller et al. 2009; Matsui et al. 2010). The expression of these genes in transgenic plants activated downstream cold-responsive genes (*COR* genes) even at warm temperature improving the cold tolerance of these plants (Liu et al. 1998; Kasuga et al. 1999). The *CBF/DREB1* pathway has been identified and described in several plant species, indicating that the CBF transcriptional cascade under the cold stress is highly conserved in the plant kingdom (Jaglo et al. 2001). A significant relation between the expression of *CBFs/DREBs* and changes in the temperature has been demonstrated, indicating that the lower the temperature the higher the expression of these genes. However, the expression can desensitize at a given low temperature (Zarka et al. 2003).

The increase in transcription levels of three *CBFs* (*CBF* 1, 2 and 3) has been observed within 15 min exposure to low temperatures. The rapid response to the cold effect strongly suggests that the 'thermometer' and signal 'transducer' of low temperature and which regulates the expressions of *CBFs* is present at warm temperatures. In this context, Gilmour et al. (1998) demonstrated that a transcription factor,

tentatively designed inducer of CBF expression (ICE), is present at warm temperature. They also reported that ICE recognizes DNA-regulatory elements in the CBF promoter and called these 'ICE boxes'. It was also reported that either 'ICE' or proteins with which it interacts are activated under the effect of low temperature and as a consequence the promoter of *CBF* genes was suggested to be activated and responsive to low temperature (Zarka et al. 2003). Chinnusamy et al. (2003) reported that *ICE1*, which is a nuclear gene encoding a MYC-like bHLH protein, has a crucial role in regulating some *CBF* genes. The role of ICE1-CBF/DREB1 singling pathway in increasing cold tolerance has been investigated in different plant species such as *A. thaliana* (Welling & Palva 2008; Chen et al. 2009; Fursova et al. 2009; Yang et al. 2011), *Cucumis sativa* (Liu et al. 2010), *Triticum aestivum* (Miller et al. 2006; Morran et al. 2011), *Hordeum vulgare* L. (Stockinger et al. 2007; Campoli et al. 2009), *Solanum tuberosum* (Behnam et al. 2007), eucalyptus (Navarro et al. 2009), *Nicotiana glauca* (Gutha & Reddy 2008), apple (Artlip et al. 2014), etc. Badawi et al. (2008) reported that there are two ICE1 homologs, *TaICE141* and *TaICE187*, in wheat and they up-regulate the expression of *CBF* group IV genes in wheat. Several studies have reported the existence of the CBF/DREB pathway and its role in increasing the cold tolerance of rice plants (Ito et al. 2006; Wang et al. 2008; Yang et al. 2009; Zhang et al. 2009; Cruz et al. 2013). Cold stress induces the expression of *OsICE1* and *OsICE2* in rice, leading to the up-regulation of *OsDREB1B*, *OSHsfA3* and *OsTPP1* (Nakamura et al. 2011). Zhao et al. (2013) indicated that the chilling tolerance of banana plant (*Musa acuminata*) was improved by jasmonate (JA) through the induction of MaMYC2. However, MaMYC2 interact with MaICE1, activating the cold signaling of the CBF-dependent pathway. Soltész et al. (2013) reported that TaCBF14 and TaCBF15 play an essential role in cold tolerance of wheat. Moreover, they reported that these genes improved the cold tolerance of transgenic plants by several degrees lower than control plants. At least five CBF genes have been identified in both apple (*Malus domestica*) and peach (*Prunus persica*) that exhibit a variety of patterns of expression in response to low temperatures (Wisniewski et al. 2014, 2015).

Fursova et al. (2009) indicated that the over-expression of *ICE2*, which is a homolog to *ICE1*, improves the cold tolerance and the expression of *CBF/DREB1B* in *Arabidopsis*. However, it was demonstrated that the hyperacetylation of histones H3 and H4 and DNA demethylation take place in the ICE-binding region of the promoter of maize (*ZmDREB1*) after cold treatment, escorted by chromatin decondensation (Hu et al. 2011). This suggests that chromatin remodeling could be one of the essential requirements for the up-regulation of *CBF/DREB1* by ICE1. However, it should be mentioned that apart from ICE proteins, there are several proteins such as LOS4, HOS1 and LOS1 proteins that have been reported to have positive effects of the induction of CBF expression (Matthew 2005).

CBF proteins, which are DNA-binding proteins, recognize the DNA-regulatory element CRT/DRE that is located in the promoter of *COR* genes. Therefore the induction of the *COR* gene expression is controlled by CBF proteins (Baker et al. 1994; Yamaguchi-Shinozaki & Shinozaki 1994). These proteins (CBF 1, 2 and 3) are encoded by three *CBF* genes located in tandem in chromosome 4 and they are considered as transcription factors having masses of about 24 kDa. These

proteins are about 90% identical in amino acid sequence and they contain a conserved DNA-binding motif of about 60 amino acids designated the AP2/ERF domain (Riechmann & Meyerowitz 1998). However, 144 AP2/ERF genes were found and described in *Arabidopsis* (Riechmann et al. 2000) and based on the comparisons of their DNA-binding domain, Sakuma et al. (2002) has classified them into 5 subgroups: AP2 subfamily (14 genes), RAV subfamily (6 genes), DREB/CBF subfamily (55 genes), ERF subfamily (65 genes) and the others which were considered as a 5th group consisting of 4 genes.

The up-regulation of *COR* (cold-regulated) genes through the activation of CRT/DRE elements in their promoters by CBF proteins is considered to be a small part of the CBF regulon since intensive studies, which have been conducted to determine the expression profiling of CBF, reported that a total of 109 genes were being assigned to CBF regulon (Fowler & Thomashow 2002; Vogel et al. 2005). However, according to the nature of proteins that are encoded, Matthew (2005) classified CBF regulon-assigned genes into four categories: the first group is the largest containing more than 50% of the proteins and these proteins have unknown function, the second group is the cryoprotective proteins which involves the *COR* proteins, the third group is the regulatory proteins and the last group is the biosynthetic proteins.

COR (cold-regulated) genes encode extremely hydrophilic proteins that are either novel or members of the dehydrins or LEA (late embryogenesis abundant) proteins (Close 2006). LEA proteins are suggested to be important for membrane stabilization and prevent protein aggregation (Hundertmark & Hincha 2008). However, most *COR* protein functional activities are still hypothetical. Artus et al. (1996) reported some evidence that the CORa protein, which is one of the most well-studied *COR* proteins, supports the stabilizing of cell membranes against freezing damage. COR15a that encodes a 15 kDa polypeptide is targeted to the chloroplast where it is processed to 9.4 kDa mature polypeptide (Lin & Thomashow 1992). Steponkus et al. (1998) reported that the COR15a polypeptide could act directly as a cryoprotective agent improving the cold tolerance of plants. Moreover, it has been demonstrated that *COR* proteins reduce the tendency of membranes forming hexagonal-II structures which is a harmful non-bilayer construction that takes place due to cellular dehydration associated with freezing (Thomashow 1999). Bravo et al. (2003) and Hara et al. (2003) have reported that *COR* proteins also have a positive role in protecting other proteins against freeze-thaw inactivation in vitro. Moreover, it has been suggested that *CORs* could play a role as a hydration buffer that isolate ions and help to re-nature unfolded proteins (Bray 1993). The essential role of *CORs* in the cold tolerance of plants was confirmed by Artus et al. (1996) who reported that over-expression of *COR15a* in transgenic *Arabidopsis* improved chloroplast freezing tolerance by about 2°C in non-acclimated plants.

In addition to *COR* genes, heat shock proteins' expression is also induced under the effect of low temperature (Timperio et al. 2008). It was also confirmed that bacterial cold shock proteins (CSPs) enhanced stress adaptation in multiple plant species by demonstrating improved stress tolerance in maize and rice. The expression of *CspA* and *CspB* in transgenic rice improved the growth of plants under the effect of a number of abiotic factors like cold, heat and water deficit. The expression of bacterial CSPs increased cold tolerance in transgenic *Arabidopsis* (Karlson & Imai 2003; Nakaminami

et al. 2006; Castiglioni et al. 2008). Some PR (pathogen-related) proteins such as PR1, PR2, PR5, PR10, PR11 and PR14 are approved to be up-regulated by low-temperature treatment (Seo et al. 2008; Zhang et al. 2010).

It should be mentioned that there is a potential connection between the cold acclimation pathway and other physiological process pathways in plants. For example, Dhillon et al. (2010) suggested a prospective relation between cold acclimation and vernalization because both the processes require an exposure to low non-freezing temperature. Moreover, the initial increase in cold tolerance observed when winter genotypes of cereals were exposed to low non-freezing was decreased with the continuous cold exposure (Fowler et al. 1996). Fowler and Limin (2004) reported that this observed decline in cold tolerance inversely parallels the accomplishment of the vernalization requirement. However, it was suggested that a controlling factor of freezing tolerance is related to a developmental alteration between the vegetative and reproductive stages and that the main vernalization gene (*VRN-1*) has an essential role in the decrease of the capacity of cold acclimation with plants' development (Limin & Fowler 2006). Stockinger et al. (2007) reported that when the winter genotype of barley carrying *vrn-1* allele is vernalized, the transcription levels of *CBF* are relatively decreased compared to non-vernalized plants. Pidal et al. (2009) reported that main differences between winter and spring cereals genotypes in terms of *vrn-1* characterization were insertions and deletions in the *vrn-1* regions located in the promoter and first intron of this gene. However, it has been reported that the main two loci which control low-temperature tolerance in wheat are the frost resistance *Fr-1* and *Fr-2* (Vágújfalvi et al. 2000). The locus of *Fr-1* was reported to be closely connected to the vernalization locus *Vrn1*, which is well known to be the main controller of the transition of the vegetative to the reproductive meristem in response to low temperature (Laudencia-Chingcuanco et al. 2011). Båga et al. (2007) reported that *Vrn-1* and *Fr-1* loci seem to be in the same locus and that they have not been genetically uncoupled. The pleiotropic properties of the *Vrn-1* locus clarify most of the related cold tolerance and winter habit (Dhillon et al. 2010). Moreover, it was reported that the *FR-2* locus contains a cluster of CBFs which are well known to have an essential role in the up-regulation of cold-induced genes (CORs) in several plant species (Campoli et al. 2009; Knox et al. 2010; Laudencia-Chingcuanco et al. 2011). Manipulating the expression of genes as response to low-temperature stress is followed by increase in metabolites, some of which are proved to have a protective role against the damage of low-temperature stress. However, plants experience changes in their pattern of gene expression and protein products when they undergo low-temperature stress and manipulating and engineering of CBF genes have shown potential practical application for breeding cold tolerance in crops and horticultural plants suitable for temperate regions (Sanghera et al. 2011).

The effect of phytohormones on plant cold tolerance

Plant hormone levels are altered under abiotic stresses and they decrease plant growth. Cold response is one of the most important environmental stresses that induces growth repression and reduced yield. Plant hormones activities induce the cascades of cold-responsive genes.

The plant hormone ABA plays an essential role as a chemical signal in response to biotic and abiotic stresses such as salt, cold, drought and wounding. ABA stimulates several changes in plant physiological, molecular and developmental progressions resulting in plant adaptation to the stress environment (Ton et al. 2009) as well as it is characterized as an intracellular messenger which has been proven to perform the critical function of the heart of signaling operation (Xue-Xuan et al. 2010). Abiotic stresses induce the synthesis of ABA which in turn activates the expression of stress-related genes and stomatal closure (Lee & Luan 2012).

ABA has an essential role in the induction of various stress signals controlling downstream stress responses. Therefore, it is necessary that plants adjust their ABA content constantly in response to changes in physiological and environmental conditions. Plants respond to stresses by two main mechanisms described as ABA-dependent and ABA-independent pathways. The expression of ABA-responsive genes could be induced by several transcription factors such as DREB2A/2B, AREB1, RD22BP1 and MYC/MYB through interacting with their corresponding cis-acting elements such as DRE/CRT, ABRE and MYCRS/MYBRS, respectively (Tuteja 2007). Koornneef et al. (1998) have reported ABA-deficient mutants namely *aba1*, *aba2* and *aba3* in *Arabidopsis*. However, while no differences were observed between the growth of these mutants and wild-type plants under no stress conditions, ABA-deficient mutants readily wilt and die under drought stress and showed poor growth under salt stress. ABA can also be essential for freezing tolerance through the induction of dehydration tolerance genes (Xiong et al. 2001).

Several ABA-induced genes such as *Arabidopsis* cold-regulated (*COR*) genes, *RAB18*, *LTI78* and *KIN2*, are up-regulated under the effect of various abiotic stresses such as cold, salt and drought (Knight et al. 2004). However, although Shinozaki and Yamaguchi-Shinozaki (2000) described several signaling pathways including both ABA-dependent and ABA-independent pathways resulting in *COR* gene (cold-responsive genes) expression up-regulation, Thomashow (1999) and Shinozaki and Yamaguchi-Shinozaki (2000) reported cold effects on the induction of gene expression was mainly through an ABA-independent pathway. The existence of an ABRE cis-acting element (ABA-responsive element) is an essential requirement for the up-regulation of ABA-induced gene expression (Shinozaki & Yamaguchi-Shinozaki 2000). It has been demonstrated using genetic analysis that there is no strong connection between ABA-dependent and ABA-independent pathways; however, the mechanism involved could usually crosstalk or sometimes converge in the signaling pathway (Uno et al. 2000). In view of this information, calcium was suggested to be a strong candidate that can mediate such crosstalks since calcium concentration was observed to rapidly increase in the plant cells under the effect of ABA and various abiotic stresses such as drought, cold and salinity (Mahajan & Tuteja 2005). Moreover, various changes in plant molecular and biochemical such as the transcript accumulation of *RD29A* gene and the accumulation of proline were observed and mediated by both ABA-dependent and ABA-independent pathways (Tuteja 2007) (Figure 3).

Finkelstein et al. (2002) reported an important role of ABA in the induction of *LEA* gene expression. Bies-Etheve et al. (2008) also reported that the application of exogenous ABA induced the expression of specific *LEA* genes in plant

vegetative tissues. The expression of *LEA* genes during the development of seed and under abiotic stress was up-regulated by ABA (Dalal et al. 2009). It has also been reported that ABA induces the expression of *LEA* genes resulting in an increase in the desiccation tolerance of cultured embryos (Leung & Giraudat 1998). The role of ABA in the up-regulation of *LEA* genes is considered to be one of the mechanisms that ABA has to increase plant drought and freezing tolerance.

Salicylic acid (SA) has been reported to be accumulated under the chilling effect in different plant species (Wan et al. 2009; Kosova et al. 2012). Moreover, the application of SA improved the cold tolerance of several plant species such as potato, rice, maize, etc. (Ahmad & Prasad 2012). Nevertheless, the over-accumulation of SA in several mutants of *Arabidopsis* caused a freezing sensitivity (Miura & Ohta 2010). It seems that treatment duration and the concentration of SA play essential roles in the response of plant species.

Gibberellin (GA) is the other plant hormone altered in plants under cold stress. It has been evident that GA plays a crucial role in the plant growth-regulatory mechanisms. It has been found that GA is involved in the expression of CRT/DRE-binding factor gene which in turn confers tolerance to drought, salt and cold stress (Niu et al. 2014). Moreover, GA is associated with SA/JA balance in the CBF-mediated stress response (Niu et al. 2014). The key components of GA is the nuclear localized proteins (DELLA) that repress plant growth; however recent advances indicated that DELLA has a role in many plant growth aspects especially those that are affected by environmental stresses (Achard et al. 2008). Achard et al. (2008) investigated the relationship between *CBF1* and GA pathways and indicated that constitutive expression of *CBF1* leads to less bioactivity of GA resulting in dwarfism in plants and delay in flowering. *RGL* genes, which code for DELLA proteins, act to restrain growth, whereas GA promotes growth by overcoming DELLA-mediated growth restraint (Claeys et al. 2014).

Plant phytohormone JA also plays an essential role as an important regulatory signal in plant cold tolerance (Cao et al. 2009; Sayyari et al. 2011; Zhao et al. 2013). It has been proved that the external application of JA significantly enhanced cold tolerance in plants with or without acclimation. Moreover, blocking of the endogenous JA increased the sensitivity to the cold stress. It has been proved that JA up-regulated the *CBF/DREB1* signaling pathway (Hu et al. 2013).

Other hormones such as ethylene and cytokinin (CK) also have an important role in cold tolerance. Although it has been found that CK content was decreased in plants exposed to stress, the next application of CK improved cold tolerance in plant. However, stress-related genes at transcriptional-level pattern have been identified by CK but the exact role of CK in cold stress is still unknown (Zwack & Rashotte 2015). Ethylene was proved to increase during low temperature in grapevine (Sun et al. 2016). Finally, phytohormones engineering is very promising in plant response to environmental stresses; however, more research is still needed in this field (Wani et al. 2016).

The role of molybdenum (Mo) in abiotic stress tolerance

Molybdenum (Mo) is an essential trace element present in the soil and is required by both plants and animals. Many

oxidation statuses of Mo ranging from zero to VI have been reported in agricultural soils. However, VI is considered to be the most existed state (Sun et al. 2009). Mo belongs to a group of trace elements that are required, by organisms in a minute amount; in spite of this, unavailability of Mo is lethal for microorganisms. Mo has to be bounded to a pterin in order to form a Mo co-factor (Mo-co) (Mendel & Bittner 2006). Mo is not biologically active but it forms a part of an organic complex called the Mo-co. Mo-co is very important for the Mo-requiring enzymes (molybdoenzymes) that have been reported in most biological systems (Williams & Frausto da Silva 2002) and all Mo-dependent enzymes (molybdoenzymes) hold a mononuclear element of Mo-co, which consists of Mo coordinated to an organic tricyclic pyranopterin moiety, referred to as molybdopterin. Mo-co in different variants represents the active compound at the catalytic site of all Mo-containing enzymes in nature (except nitrogenase), that is several enzymes catalyzing divers keys oxotransfer reactions in the global carbon, sulfur and nitrogen metabolism (Mendel & Bittner 2006). Furthermore, even if Mo is available in the medium, it has to be complexed by a special co-factor to confer catalytic activity to several enzymes. Mo is an important element for more than 40 enzymes, 4 of which have been found in plants (Mendel & Hansch 2002) and include nitrate reductase (NR) which is involved in nitrogen fixation and assimilation, xanthine dehydrogenase/oxidase (XDH) which plays an important role in purine catabolism, aldehyde oxidase (AO) that plays an essential part in the synthesis of indole-3 acetic acid and ABA and sulfite oxidase (SO) which have a role in sulfur metabolism (Kaiser et al. 2005). XDH and AO contain a monoxo-Mo-co which needs Mo-co insertion and then consequent sulfuration of the Mo center to activate the Mo-co protein complex. SO and NR have a dioxo-Mo-co which motivates the protein when it is inserted into the protein complex (Mendel & Hansch 2002).

Li et al. (2001) proposed that Mo is involved in the amelioration of wheat frost tolerance. Several mechanisms, of which Mo can improve the frost tolerance, have been suggested. However, the mechanism for this enhancement has not been definitively determined. Mo increases the activity of the anti-oxidative enzymes SOD, CAT and POX resulting in an increase in plant anti-oxidative defenses under low-temperature stress (Yu et al. 1999) since frost stress leads to the production of ROS that has cellular damage effects on plant tissues (Sattler et al. 2000). Sun et al. (2006) demonstrated that Mo application increases the capacity of scavenging the ROS resulting in the reduction in membrane damage of winter wheat under low temperature.

Another hypothesis proposed for the role of Mo in plant cold tolerance was that Mo increases the activity of AO which has an important role in the production of ABA. ABA triggers bZIP and the up-regulation of the ABA-dependent COR gene expression pathways (Sun et al. 2009) and thereby protects against frost damage. However, the role of Mo in the biosynthesis of ABA suggests a significant role of this element in plant drought and salinity tolerance.

A third mechanism is suggested to be through NR and enhancing nitrate assimilation since Mo has significantly increased NR activity in wheat as it has been found that NR activity significantly increased in Mo-treated wheat and was associated with protein accumulation (Yu et al. 1999; Hale et al. 2001; Hamdia et al. 2005). Al-Issawi et al. (2013) recently reported a fourth mechanism of Mo effects on the




Figure 3. Cold-regulated gene induction pathways.

plant cold tolerance through the up-regulation of the CBF pathway which in turns lead to *COR* up-regulation and improvements in frost tolerance.

Utilizing the infrared camera in cold tolerance in planta

Many economic inputs are used to provide considerable protection against freezing events in plants or parts of plants, for example, rowcovers, windmills, smudge pots or spraying water on plants. All these methods seem to be ineffective in giving the required protection to the plants; however very little is known about the mechanisms associated with freezing of plants in the field conditions; therefore this gave a chance to use the most effective method of frost protection which is the use of high resolution of infrared camera directly in the field (Wisniewski et al. 2008). Thermal infrared imaging techniques are very valuable as an aid in understanding the physiological effects of frost on plants and can be used both in the field and under artificial conditions. This technique is based on measuring the heat released when different fractions of water freeze within tissue (Quamme et al. 1972; Ketchie & Kammereck 1987; Fuller & Wisniewski 1998). The thermal infrared camera produces an image in a way exactly analogous to forming images from visible light with familiar electronic imaging devices like video or digital cameras. Thermal cameras contain an array of detectors that convert infrared radiation (thermal energy) to electrical output signals whose amplitude is proportional to the amount of the energy detected. These output signals can be processed as a normal video image in gray-scale image of the scene. The gray-scale value in the images represents the amount of thermal energy detected from a given point in view and is proportional to the radiant temperature of that point. Hotter radiant temperatures are normally displayed as whiter tones in an image, and colder temperature as blacker tones (Colwell 1983). More recently, gray-scale images can be converted to color-enhanced images using computing software generating a range of color palettes. Since freezing events are temporal it is necessary to record the images over time either as video

images or digital video equivalents. The determination of the exact location of initial ice formation (point of ice nucleation) and how it progresses throughout the plant (ice spread) is very important information that needs to be known prior to devising methods to protect plants. Wisniewski et al. (1997) were among the first to use infrared thermal imaging technology to determine where ice first begins to form in a plant and at what temperature, based on the principle of release of heat when ice is formed (exotherm which is the process that describes the chemical reaction when energy is released usually in the form of heat). By using infrared video thermography, these researchers observed a plant freezing and followed the progress of ice formation throughout the plant. Subsequently, ice formation has been tracked by using an infrared camera system in many species including apple (*Malus ssp*), peach (*P. persica*) and pear (*Pyrus ssp*) trees and bean (*Vicia faba* L.), potato (*S. tuberosum*), strawberry (*Fragaria ssp*) and wheat plants (*T. aestivum* L.) (Wisniewski et al. 1997; Fuller & Wisniewski 1998). Computer software has also been used to analyze recordings of infrared video tape to study all stages of ice formation and to quantify the magnitude of the exotherms observed and their duration (Al-Swedi et al. 2013). Furthermore, the speed of ice front travel can also be determined by determining the start and the end times of the whole tissue being frozen and barriers to ice spread being detected (Wisniewski et al. 1997; Pearce & Fuller 2001) (Figure 4).

Of significance have been the observations that stem nodes hindered ice travel but ice forms quickly along the stem of the plant; but slows as it approaches the nodes; also, while shoots freeze quickly, ice formation progresses very slowly into the roots. In peach trees the stem was frozen before flowers and ice does not automatically form in the flowers, when shoots have frozen. Such differential freezing in various plant structures can begin to answer field observations of differential frost damage following natural freezing events.

Since the discovery that some bacteria (e.g. *Pseudomonas syringae*) can stimulate ice nucleation these have been used in freezing experiments to aid ice formation during infrared observation experiments (Wisniewski et al. 1997; Fuller &




Figure 4. The use of IR thermography technique showing the pattern of ice nucleation events in wheat spikes of non-sprayed and non-acclimated plants. The plants were in a frost chamber and an IR camera was fixed in the front window of it while plants were subjected to a freezing temperature down to -6°C at a freezing rate of 2°C down each 2 hours. (Circles refer to the start points of ice nucleation and the arrows refer to its progress pattern) (Al-Issawi 2013).

Wisniewski 1998; Pearce & Fuller 2001). Typically, plants, which were treated with droplets of INA bacteria began to freeze first compared with samples or plants treated with water droplets only. It was observed that the drop of water on the surface of a leaf froze independently, at least 2.5 minutes after the part of leaf containing the bacteria (Wisniewski et al. 1997). Fuller et al. (2007) have reported that the ice nucleation temperature was uniform with *P. syringae* but was lower in the presence of distilled water. The warmest temperatures that INA bacteria are active at -2°C whereas

distilled water droplets can supercool significantly below this. Thus it would seem that in nature, plants will always supercool to some extent and it follows that eliminating INA bacterial cells from the plant tissue surfaces allows plants to protect themselves from freezing by enabling supercooling. This approach is being used in California to protect blossom in almond and peach orchards from spring radiation frosts (Lindow et al. 1982) but has not been used systematically elsewhere. Fuller et al. (2007) have demonstrated in vitro that the use of compounds that help shed surface water decreases the




Figure 5. Avoiding ice nucleation by the supercooling mechanism in wheat (temperature dropped down to -11°C and no ice nucleation events were experienced) (Al-Issawi 2013).

incidence of dew and thereby increases the likelihood of supercooling and frost escape in potatoes, tomatoes and grapevine but these compounds have not been tested extensively in the field.

Infrared thermal imaging has been used by many researchers who have observed that while supercooling is a mechanism for avoiding frost damage (Figure 5), it does carry with it an inherent risk if the temperature drops too low and is followed by freezing.

Ice crystals that develop after substantial supercooling have the potential to cause more injury than ice crystals that form at higher temperature (Choi et al. 1999). Slowly formed ice crystals are smaller than those formed more rapidly and in addition, when ice forms slowly, water has time to escape from plant cells so that ice grows in the inter-cellular spaces and intracellular ice formation, which is lethal, is less probable.

Conclusion

The physiological and molecular mechanisms of plant cold tolerance were reviewed in this study. The important role of cold acclimation and the biochemical and molecular changes that take place through this process were revised. The regulation of cold-regulated genes (CORs) and the pathways that control this regulation were discussed. Moreover, the essential part of *CBF* genes in the up-regulation and the production of COR proteins was reviewed. Recent research has demonstrated the positive role of Mo in plant cold tolerance. The role of this element in the up-regulation of COR genes through both ABA-dependent and -independent pathways was revised. Moreover, the effects of ABA on the abiotic stress tolerance generally and on the cold tolerance especially were revised. However, it is clear that plants with less up-regulation of *CBF* gene transcripts have less acclimation capability and intolerance. However, the nature and process of this effect were discussed in this review. The use of infrared camera as a tool to improve the understanding level of ice formation in plant tissues was explained. This study provides a comprehensive overview of the recent achievement in the field of plant cold tolerance.

Disclosure statement

No potential conflict of interest was reported by the authors.

References

- Achard P, Gong F, Cheminant S, Alioua M, Hedden P, Genschik P. 2008. The cold-inducible CBF1 factor-dependent signaling pathway modulates the accumulation of the growth-repressing DELLA proteins via its effect on gibberellin metabolism. *Plant Cell*. 20:2117–2129.
- Ahmad P, Prasad MMNV. 2012. Abiotic stress responses in plants. New York: Springer.
- Al-Issawi M. 2013. The characterisation of the freezing damage response during flowering in European and Middle Eastern wheat cultivars (*Triticum aestivum* L.). [PhD]. University of Plymouth.
- Al-Issawi M, Rihan HZ, Woldie WA, Burchett S, Fuller MP. 2013. Exogenous application of molybdenum affects the expression of CBF14 and the development of frost tolerance in wheat. *Plant Physiol Biochem*. 63:77–81.
- Al-Swedi F, Rihan HZ, Al Shamar M, Lane S, Fuller MP. 2013. Genetic transformation of cauliflower with ascorbate peroxidase (APX) gene. *Acta Hortic*. Under Review.
- Artlip TS, Wisniewski ME, Norelli JL. 2014. Field evaluation of apple overexpressing a peach CBF gene confirms its effect on cold hardiness, dormancy, and growth. *Environ Exp Bot*. 106:79–86.
- Artus NN, Uemura M, Steponkus PL, Gilmour SJ, Lin CT, Thomashow MF. 1996. Constitutive expression of the cold-regulated *Arabidopsis thaliana* COR15a gene affects both chloroplast and protoplast freezing tolerance. *Proc Natl Acad Sci USA*. 93:13404–13409.
- Aryal B, Neuner G. 2010. Leaf wettability decreases along an extreme altitudinal gradient. *Oecologia*. 162:1–9.
- Badawi M, Danyluk J, Boucho B, Houde M, Sarhan F. 2007. The CBF gene family in hexaploid wheat and its relationship to the phylogenetic complexity of cereal CBFs. *Mol Genet Genomics*. 277:533–554.
- Badawi M, Reddy YV, Agharbaoui Z, Tominaga Y, Danyluk J, Sarhan F, Houde M. 2008. Structure and functional analysis of wheat ICE (inducer of CBF expression) genes. *Plant Cell Physiol*. 49:1237–1249.
- Baker SS, Wilhelm KS, Thomashow MF. 1994. The 5'-region of *Arabidopsis thaliana* cor15a has cis-acting elements that confer cold-, drought- and ABA-regulated gene expression. *Plant Mol Biol*. 24:701–713.
- Behnam B, Kikuchi A, Celebi-Toprak F, Kasuga M, Yamaguchi-Shinozaki K, Watanabe KN. 2007. *Arabidopsis* rd29A: DREB1A enhances freezing tolerance in transgenic potato. *Plant Cell Rep*. 26:1275–1282.
- Bies-Etheve N, Gaubier-Comella P, Debures A, Lasserre E, Jobet E, Raynal M, Cooke R, Delseny M. 2008. Inventory, evolution and expression profiling diversity of the LEA (late embryogenesis abundant) protein gene family in *Arabidopsis thaliana*. *Plant Mol Biol*. 67:107–124.
- Bravo LA, Gallardo J, Navarrete A, Olave N, Martínez J, Alberdi M, Close TJ, Corcuera LJ. 2003. Cryoprotective activity of a cold-induced dehydrin purified from barley. *Physiol Plant*. 118:262–269.
- Bray EA. 1993. Molecular responses to water deficit. *Plant Physiol*. 103:1035–1040.
- Bray EA. 2004. Genes commonly regulated by water-deficit stress in *Arabidopsis thaliana*. *J Exp Bot*. 55:2331–2341.
- Bäga M, Chodaparambil SV, Limin AE, Pecar M, Fowler DB, Chibbar RN. 2007. Identification of quantitative trait loci and associated candidate genes for low-temperature tolerance in cold-hardy winter wheat. *Funct Integr Genomics*. 7:53–68.
- Campoli C, Matus-Cádiz M, Pozniak C, Cattivelli L, Fowler DB. 2009. Comparative expression of Cbf genes in the Triticeae under different acclimation induction temperatures. *Mol Genet Genomics*. 282:141–152.
- Cao S, Zheng Y, Wang K, Jin P, Rui H. 2009. Methyl jasmonate reduces chilling injury and enhances antioxidant enzyme activity in postharvest loquat fruit. *Food Chem*. 115:1458–1463.
- Castiglioni P, Warner D, Bensen RJ, Anstrom DC, Harrison J, Stoecker M, Abad M, Kumar G, Salvador S, D'ordine R, et al. 2008. Bacterial RNA chaperones confer abiotic stress tolerance in plants and improved grain yield in maize under water-limited conditions. *Plant Physiol*. 147:446–455.
- Chen HH, Brenner ML, Li PH. 1993. Involvement of abscisic acid in potato cold acclimation. *Plant Physiol*. 71:362–365.
- Chen M, Xu Z, Xia L, Li L, Cheng X, Dong J, Wang Q, Ma Y. 2009. Cold-induced modulation and functional analyses of the DRE-binding transcription factor gene, GmDREB3, in soybean (*Glycine max* L.). *J Exp Bot*. 60:121–135.
- Chen L, Zhang Y, Ren Y, Xu J, Zhang Z, Wang Y. 2012. Genome-wide identification of cold-responsive and new microRNAs in *Populus tomentosa* by high-throughput sequencing. *Biochem Biophys Res Commun*. 417:892–896.
- Chinnusamy V, Ohta M, Kanrar S, Lee B, Hong X, Agarwal M, Zhu JK. 2003. ICE1: a regulator of cold-induced transcriptome and freezing tolerance in *Arabidopsis*. *Gene Dev*. 17:1043–1054.
- Chinnusamy V, Zhu J-K, Sunkar R. 2010. Plant stress tolerance: Methods and Protocols. Gene regulation during cold stress acclimation in plants. Springer; p. 39–55.
- Choi D-W, Zhu B, Close T. 1999. The barley (*Hordeum vulgare* L.) dehydrin multigene family: sequences, allele types, chromosome assignments, and expression characteristics of 11 Dhn genes of cv Dicktoo. *Theor Appl Genet*. 98:1234–1247.
- Claeys H, De Bodt S, Inzé D. 2014. Gibberellins and DELLAs: central nodes in growth regulatory networks. *Trends Plant Sci*. 19:231–239.
- Close TJ. 2006. Dehydrins: a commonality in the response of plants to dehydration and low temperature. *Physiol Plant*. 100:291–296.

- Colwell RN. 1983. Manual of remote sensing. Vol. 1. 2nd ed. Falls Church (VA): American Society of Photogrammetry; p. 1232.
- Cruz RPD, Sperotto RA, Cargnelutti D, Adamski JM, Freitasterra T, Fett JP. 2013. Avoiding damage and achieving cold tolerance in rice plants. *Food Energy Secur.* 2:96–119.
- Cushman JC, Bohnert HJ. 2000. Genomic approaches to plant stress tolerance. *Curr Opin Plant Biol.* 3:117–124.
- Dahal K, Kane K, Gadapati W, Webb E, Savitch LV, Singh J, Sharma P, Sarhan F, Longstaffe FJ, Grodzinski B. 2012. The effects of phenotypic plasticity on photosynthetic performance in winter rye, winter wheat and *Brassica napus*. *Physiol Plant.* 144:169–188.
- Dalal M, Tayal D, Chinnusamy V, Bansal KC. 2009. Abiotic stress and ABA-inducible Group 4 *LEA* from *Brassica napus* plays a key role in salt and drought tolerance. *J Biotechnol.* 139:137–145.
- Deane C. 1994. Whole plant and tissue culture studies on frost tolerance in cauliflower [PhD]. The National University of Ireland.
- Denesik TJ. 2007. Quantitative expression analysis of four low-temperature-tolerance-associated genes during cold acclimation in wheat (*Triticum aestivum* L.) [Master of Science]. University of Saskatchewan.
- Dhillon T, Pearce SP, Stockinger EJ, Distelfeld A, Li C, Knox AK, Vashegyi I, Vágújfalvi A, Galiba G, Dubcovsky J. 2010. Regulation of freezing tolerance and flowering in temperate cereals: the VRN-1 connection. *Plant Physiol.* 153:1846–1858.
- Doherty CJ, Van Buskirk HA, Myers SJ, Thomashow MF. 2009. Roles for *Arabidopsis* CAMTA transcription factors in cold-regulated gene expression and freezing tolerance. *Plant Cell Online.* 21:972–984.
- Dörffling K, Dörffling H, Lesselich G, Luck E, Zimmermann C, Melz G, Jürgens HU. 1997. Heritable improvement of frost tolerance in winter wheat by in vitro-selection of hydroxyproline-resistant proline over-producing mutants. *Euphytica.* 93:1–10.
- Eckardt NA. 2009. CAMTA proteins: a direct link between calcium signals and cold acclimation? *Plant Cell Online.* 21:697–697.
- Finkelstein RR, Gampala SSL, Rock CD. 2002. Absciscic acid signaling in seeds and seedlings. *Plant Cell Online.* 14:S15–S45.
- Fowler D, Chauvin L, Limin A, Sarhan F. 1996. The regulatory role of vernalization in the expression of low-temperature-induced genes in wheat and rye. *Theor Appl Genet.* 93:554–559.
- Fowler D, Limin A. 2004. Interactions among factors regulating phenological development and acclimation rate determine low-temperature tolerance in wheat. *Ann Bot.* 94:717–724.
- Fowler S, Thomashow MF. 2002. *Arabidopsis* transcriptome profiling indicates that multiple regulatory pathways are activated during cold acclimation in addition to the CBF cold response pathway. *Plant Cell.* 14:1675–1690.
- Fuller PM, Fuller AM, Spyridon K, Jack C, Troy F. 2007. The freezing characteristics of wheat at ear emergence. *Eur J Agron.* 26:435–441.
- Fuller MP, Wisniewski M. 1998. The use of infrared thermal imaging in the study of ice nucleation and freezing of plants. *J Therm Biol.* 23:81–89.
- Fursova OV, Pogorelko GV, Tarasov VA. 2009. Identification of ICE2, gene involved in cold acclimation which determines freezing tolerance in *Arabidopsis thaliana*. *Gene.* 429:98–103.
- Gao JP, Chao DY, Lin HX. 2007. Understanding abiotic stress tolerance mechanisms: recent studies on stress response in rice. *J Integr Plant Biol.* 49:742–750.
- George MF, Becwar MR, Burke MJ. 1982. Freezing avoidance by deep super-cooling of tissue water in winter-hardy plants. *Cryobiology.* 19:628–639.
- Gery C, Zuther E, Schulz E, Legoupi J, Chauveau A, Mckhann H, Hinch DK, Téoulé E. 2011. Natural variation in the freezing tolerance of *Arabidopsis thaliana*: effects of RNAi-induced CBF depletion and QTL localisation vary among accessions. *Plant Sci.* 180:12–23.
- Ghildiyal M, Zamore PD. 2009. Small silencing RNAs: an expanding universe. *Nat Rev Genet.* 10:94–108.
- Gilmour SJ, Zarka DG, Stockinger EJ, Salazar MP, Houghton JM, Thomashow MF. 1998. Low temperature regulation of the *Arabidopsis* CBF family of AP2 transcriptional activators as an early step in cold-induced COR gene expression. *Plant J.* 16:433–442.
- Gusta LV, Wisniewski M. 2013. Understanding plant cold hardiness: an opinion. *Physiol Plant.* 147:4–14.
- Gusta LV, Wisniewski ME, Tanino KK. 2009. Plant cold hardiness: from the laboratory to the field. UK: CAB.
- Gutha LR, Reddy AR. 2008. Rice DREB1B promoter shows distinct stress-specific responses, and the overexpression of cDNA in tobacco confers improved abiotic and biotic stress tolerance. *Plant Mol Biol.* 68:533–555.
- Guy CL. 1990. Cold acclimation and freezing stress tolerance: role of protein metabolism. *Annu Rev Plant Biol.* 41:187–223.
- Guy C, Haskell D, Li QB. 1998. Association of proteins with the stress 70 molecular chaperones at low temperature: evidence for the existence of cold labile proteins in spinach. *Cryobiology.* 36:301–314.
- Hacker J, Ladinig U, Wagner J, Neuner G. 2011. Inflorescences of alpine cushion plants freeze autonomously and may survive subzero temperatures by supercooling. *Plant Sci.* 180:149–156.
- Hale KL, Mcgrath SP, Lombi E, Stack SM, Terry N, Pickering IJ, George GN, Pilon-Smits EAH. 2001. Molybdenum sequestration in *Brassica* species. A role for anthocyanins. *Plant Physiol.* 126:1391–1402.
- Hamdia HM, El-Komy HM, Shaddad MAK, Hetta AM. 2005. Effect of molybdenum of nitrogenase activities of wheat inoculated with *Azospirillum brasilense* growth under drought stress. *Gen Appl Plant Physiol.* 31:43–54.
- Hannah MA, Heyer AG, Hinch DK. 2005. A global survey of gene regulation during cold acclimation in *Arabidopsis thaliana*. *PLoS Genet.* 1:e26.
- Hara M, Terashima S, Fukaya T, Kuboi T. 2003. Enhancement of cold tolerance and inhibition of lipid peroxidation by citrus dehydrin in transgenic tobacco. *Planta.* 217:290–298.
- Heidarvand L, Maali Amiri R. 2010. What happens in plant molecular responses to cold stress? *Acta Physiol Plant.* 32:419–431.
- Hu Y, Jiang L, Wang F, Yu D. 2013. Jasmonate regulates the inducer of CBF expression—c-repeat binding factor/DRE binding factor1 cascade and freezing tolerance in *Arabidopsis*. *Plant Cell.* 25:2907–2924.
- Hundertmark M, Hinch DK. 2008. LEA (late embryogenesis abundant) proteins and their encoding genes in *Arabidopsis thaliana*. *BMC Genomics.* 9:118.
- Hu Y, Zhang L, Zhao L, Li J, He S, Zhou K, Yang F, Huang M, Jiang L, Li L. 2011. Trichostatin A selectively suppresses the cold-induced transcription of the ZmDREB1 gene in maize. *PLoS ONE.* 6:e22132.
- Ito Y, Katsura K, Maruyama K, Taji T, Kobayashi M, Seki M, Shinozaki K, Yamaguchi-Shinozaki K. 2006. Functional analysis of rice DREB1/CBF-type transcription factors involved in cold-responsive gene expression in transgenic rice. *Plant Cell Physiol.* 47:141–153.
- Ivanov A, Rosso D, Savitch L, Stachula P, Rosembert M, Oquist G, Hurry V, Hüner N. 2012. Implications of alternative electron sinks in increased resistance of PSII and PSI photochemistry to high light stress in cold-acclimated *Arabidopsis thaliana*. *Photosynth Res.* 113:191–206.
- Jaglo KR, Kleff S, Amundsen KL, Zhang X, Haake V, Zhang JZ, Deits T, Thomashow MF. 2001. Components of the *Arabidopsis* C-repeat/dehydration-responsive element binding factor cold-response pathway are conserved in *Brassica napus* and other plant species. *Plant Physiol.* 127:910–917.
- Jan N, Mahboob UI H, Andrabi KI. 2009. Cold resistance in plants: a mystery unresolved. *Electron J Biotechnol.* 12:14–15.
- Jenks MA, Wood AJ. 2010. Genes for plant abiotic stress. Ames (IA): Wiley-Blackwell.
- Kaiser BN, Gridley KL, Brady JN, Phillips T, Tyerman SD. 2005. The role of molybdenum in agricultural plant production. *Ann Bot.* 96:745–754.
- Karlson D, Imai R. 2003. Conservation of the cold shock protein of *Escherichia coli*. *Plant Physiol.* 131:12–15.
- Kasuga M, Liu Q, Miura S, Yamaguchi-Shinozaki K, Shinozaki K. 1999. Improving plant drought, salt, and freezing tolerance by gene transfer of a single stress-inducible transcription factor. *Nat Biotechnol.* 17:287–291.
- Ketchie DO, Kammereck R. 1987. Seasonal variation of cold resistance in *Malus* woody tissue as determined by differential thermal analysis and viability tests. *Can J Bot.* 65:2640–2645.
- Kishitani S, Watanabe K, Yasuda S, Arakawa K, Takabe T. 1994. Accumulation of glycine betaine during cold acclimation and freezing tolerance in leaves of winter and spring barley plants. *Plant Cell Environ.* 17:89–95.
- Knight H, Knight MR. 2000. Imaging spatial and cellular characteristics of low temperature calcium signature after cold acclimation in *Arabidopsis*. *J Exp Bot.* 51:1679–1686.
- Knight H, Zarka DG, Okamoto H, Thomashow ME, Knight MR. 2004. Absciscic acid induces CBF gene transcription and subsequent induction of cold-regulated genes via the CRT promoter element. *Plant Physiol.* 135:1710–1717.

- Knox A, Dhillon T, Cheng H, Tondelli A, Pecchioni N, Stockinger E. 2010. CBF gene copy number variation at Frost Resistance-2 is associated with levels of freezing tolerance in temperate-climate cereals. *Theor Appl Genet*. 121:21–35.
- Kocsy G, Athmer B, Perovic D, Himmelbach A, Szűcs A, Vashegyi I, Schweizer P, Galiba G, Stein N. 2010. Regulation of gene expression by chromosome 5A during cold hardening in wheat. *Mol Genet Genomics*. 283:351–363.
- Kocsy G, Galiba G, Brunold C. 2008. Role of glutathione in adaptation and signalling during chilling and cold acclimation in plants. *Physiol Plant*. 113:158–164.
- Koornneef M, Léon-Kloosterziel KM, Schwartz SH, Zeevaart JAD. 1998. The genetic and molecular dissection of abscisic acid biosynthesis and signal transduction in *Arabidopsis*. *Plant Physiol Biochem*. 36:83–89.
- Kosová K, Prášil IT, Vítámvás P, Dobrev P, Motyka V, Floková K, Novák O, Turečková V, Rolčík J, Pešek B. 2012. Complex phytohormone responses during the cold acclimation of two wheat cultivars differing in cold tolerance, winter Samanta and spring Sandra. *J Plant Physiol*. 169:567–576.
- Koster KL, Lynch DV. 1992. Solute acclimation and compartmentation during the cold-acclimation of Puma Rye. *Plant Physiol*. 98:108–113.
- Kreps JA, Wu YJ, Chang HS, Zhu T, Wang X, Harper JF. 2002. Transcriptome changes for *Arabidopsis* in response to salt, osmotic, and cold stress. *Plant Physiol*. 130:2129–2141.
- Kwak JM, Mori IC, Pei Z-M, Leonhardt N, Torres MA, Dangel JL, Bloom RE, Bodde S, Jones JDG, Schroeder JI. 2003. NADPH oxidase AtrbohD and AtrbohF genes function in ROS-dependent ABA signaling in *Arabidopsis*. *EMBO J*. 22:2623–2633.
- Laudencia-Chingcuanco D, Ganeshan S, You F, Fowler B, Chibbar R, Anderson O. 2011. Genome-wide gene expression analysis supports a developmental model of low temperature tolerance gene regulation in wheat (*Triticum aestivum* L.). *BMC Genomics*. 12:299.
- Lee SC, Luan S. 2012. ABA signal transduction at the crossroad of biotic and abiotic stress responses. *Plant Cell Environ*. 35:53–60.
- Leung J, Giraudat J. 1998. Abscisic acid signal transduction. *Annu Rev Plant Physiol Plant Mol Biol*. 49:199–222.
- Levitt J. 1980. Responses of plants to environmental stress. Chilling, freezing, and high temperature stresses. New York: Academic Press.
- Lichtenthaler HK. 2006. The stress concept in plants: an introduction. *Ann NY Acad Sci*. 851:187–198.
- Limin AE, Fowler DB. 2006. Low-temperature tolerance and genetic potential in wheat (*Triticum aestivum* L.): response to photoperiod, vernalization, and plant development. *Planta*. 224:360–366.
- Lindow SE, Arny DC, Upper CD. 1982. Bacterial ice nucleation: a factor in frost injury to plants. *Plant Physiol*. 70:1084–1089.
- Lin C, Thomashow MF. 1992. DNA sequence analysis of a complementary DNA for cold-regulated *Arabidopsis* gene cor15 and characterization of the COR 15 polypeptide. *Plant Physiol*. 99:519–525.
- Li C, Rudi H, Stockinger E, Cheng H, Cao M, Fox S, Mockler T, Westereng B, Fjellheim S, Rognli O. 2012. Comparative analyses reveal potential uses of *Brachypodium distachyon* as a model for cold stress responses in temperate grasses. *BMC Plant Biol*. 12:65.
- Lissarre M, Ohta M, Sato A, Miura K. 2010. Cold-responsive gene regulation during cold acclimation in plants. *Plant Signal Behav*. 5:948–952.
- Liu L, Duan L, Zhang J, Zhang Z, Mi G, Ren H. 2010. Cucumber *Cucumis sativus* L. over-expressing cold-induced transcriptome regulator *ICE1* exhibits changed morphological characters and enhances chilling tolerance. *Sci Hortic*. 124:29–33.
- Liu Q, Kasuga M, Sakuma Y, Abe H, Miura S, Yamaguchi-Shinozaki K, Shinozaki K. 1998. Two transcription factors, DREB1 and DREB2, with an EREBP/AP2 DNA binding domain separate two cellular signal transduction pathways in drought- and low-temperature-responsive gene expression, respectively, in *Arabidopsis*. *Plant Cell*. 10:1391–1406.
- Li W, Wang Z, Mi G, Han X, Zhang F. 2001. Molybdenum deficiency in winter wheat seedlings as enhanced by freezing temperature. *J Plant Nutr*. 24:1195–1203.
- Lv D-K, Bai X, Li Y, Ding X-D, Ge Y, Cai H, Ji W, Wu N, Zhu Y-M. 2010. Profiling of cold-stress-responsive miRNAs in rice by microarrays. *Gene*. 459:39–47.
- Lynch DV, Steponkus PL. 1987. Plasma-membrane lipid alternation associated with cold-acclimation of winter rye seedlings (*Secale cereale* cv Puma). *Plant Physiol*. 83:761–767.
- Mahajan S, Tuteja N. 2005. Cold, salinity and drought stresses: an overview. *Arch Biochem Biophys*. 444:139–158.
- Mahmood R. 2002. In vitro effect of salt on the vigor of potato (*Solanum tuberosum* L.) plantlets. *Int J Biotechnol*. 1:73–77.
- Marc RK, Alexander WS, Helen JR, Paul ML. 2010. Calcium-regulation of cold gene expression. Society for Experimental Biology Annual meeting. Prague: Portland Press; p. 306.
- Maruyama K, Sakuma Y, Kasuga M, Ito Y, Seki M, Goda H, Shimada Y, Yoshida S, Shinozaki K, Yamaguchi-Shinozaki K. 2004. Identification of cold-inducible downstream genes of the *Arabidopsis* DREB1A/CBF3 transcriptional factor using two microarray systems. *Plant J*. 38:982–993.
- Matsui A, Ishida J, Morosawa T, Okamoto M, Kim J-M, Kurihara Y, Kawashima M, Tanaka M, To TK, Nakaminami K. 2010. *Arabidopsis* tiling array analysis to identify the stress-responsive genes. *Plant stress tolerance*. Springer.
- Matthew AJ. 2005. Plant abiotic stress. (IN): Blackwell.
- Mauger JP. 2012. Role of the nuclear envelope in calcium signalling. *Biol Cell*. 104:70–83.
- Mazars C, Brière C, Bourque S, Thuleau P. 2011. Nuclear calcium signaling: an emerging topic in plants. *Biochimie*. 93:2068–2074.
- Mckersie BD, Bowley SR. 1997. Active oxygen and freezing tolerance in transgenic plants. *Plant cold hardiness*. New York (NY): Plenum Press; p. 203–214.
- Mckhann H, Gery C, Berard A, Leveque S, Zuther E, Hinch D, De Mita S, Brunel D, Teoule E. 2008. Natural variation in CBF gene sequence, gene expression and freezing tolerance in the Versailles core collection of *Arabidopsis thaliana*. *BMC Plant Biol*. 8:105.
- Medina J, Catalá R, Salinas J. 2011. The CBFs: three *Arabidopsis* transcription factors to cold acclimate. *Plant Sci*. 180:3–11.
- Mendel RR, Bittner F. 2006. Cell biology of molybdenum. *Biochim Biophys Acta (BBA) – Mol Cell Res*. 1763:621–635.
- Mendel RR, Hansch R. 2002. Molybdoenzymes and molybdenum cofactor in plants. *J Exp Bot*. 53:1689–1698.
- Miller A, Galiba G, Dubcovsky J. 2006. A cluster of 11 CBF transcription factors is located at the frost tolerance locus Fr-A m 2 in *Triticum monococcum*. *Mol Genet Genomics*. 275:193–203.
- Minami A, Fujiwara M, Furuto A, Fukao Y, Yamashita T, Kamo M, Kawamura Y, Uemura M. 2009. Alterations in detergent-resistant plasma membrane microdomains in *Arabidopsis thaliana* during cold acclimation. *Plant Cell Physiol*. 50:341–359.
- Miura K, Furumoto T. 2013. Cold signaling and cold response in plants. *Int J Mol Sci*. 14:5312–5337.
- Miura K, Ohta M. 2010. SIZ1, a small ubiquitin-related modifier ligase, controls cold signaling through regulation of salicylic acid accumulation. *J Plant Physiol*. 167:555–560.
- Mizoi J, Shinozaki K, Yamaguchi-Shinozaki K. 2012. AP2/ERF family transcription factors in plant abiotic stress responses. *Biochim Biophys Acta*. 1819:86–96.
- Monroe J, McGovern C, Lasky JR, Grogan K, Beck J, McKay JK. 2016. Adaptation to warmer climates by parallel functional evolution of CBF genes in *Arabidopsis thaliana*. *Mol Ecol*. 25:3632–3644.
- Monroy AF, Sangwan V, Dhindsa RS. 1998. Low temperature signal transduction during cold acclimation: protein phosphatase 2A as an early target for cold-inactivation. *Plant J*. 13:653–660.
- Morran S, Eini O, Pyvovarenko T, Parent B, Singh R, Ismagul A, Eliby S, Shirley N, Langridge P, Lopato S. 2011. Improvement of stress tolerance of wheat and barley by modulation of expression of DREB/CBF factors. *Plant Biotechnol J*. 9:230–249.
- Murelli C, Rizza F, Albini FM, Dulio A, Terzi V, Cattivelli L. 1995. Metabolic changes associated with cold-acclimation in contrasting cultivars of barley. *Physiol Plant*. 94:87–93.
- Nakaminami K, Karlson DT, Imai R. 2006. Functional conservation of cold shock domains in bacteria and higher plants. *Proc Natl Acad Sci USA*. 103:10122–10127.
- Nakamura J, Yuasa T, Huang TT, Harano K, Tanaka S, Iwata T, Phan T, Iwaya-Inoue M. 2011. Rice homologs of inducer of CBF expression (OsICE) are involved in cold acclimation. *Plant Biotechnol*. 28:303–309.
- Navarro M, Marque G, Ayax C, Keller G, Borges JP, Marque C, Teulière C. 2009. Complementary regulation of four Eucalyptus CBF genes under various cold conditions. *J Exp Bot*. 60:2713–2724.
- Niu S, Gao Q, Li Z, Chen X, Li W. 2014. The role of gibberellin in the CBF1-Mediated Stress-Response Pathway. *Plant Mol Biol Rep*. 32:852–863.

- Nomura M, Muramoto Y, Yasuda S, Takabe T, Kishitani S. 1995. The acclimation of glycinebetaine during cold-acclimation early and late cultivars of barley. *Euphytica*. 83:247–250.
- Norén L, Kindgren P, Stachula P, Rühl M, Eriksson ME, Hurry V, Strand Å. 2016. Circadian and plastid signaling pathways are integrated to ensure correct expression of the CBF and COR genes during photo-periodic growth. *Plant Physiol*. 171:1392–1406.
- Pearce RS. 2001. Plant freezing and damage. *Ann Bot*. 87:417–424.
- Pearce RS, Fuller MP. 2001. Freezing of barley studied by infrared video thermography. *Plant Physiol*. 125:227–240.
- Pearce S, Zhu J, Boldizsár Á, Vágújfalvi A, Burke A, Garland-Campbell K, Galiba G, Dubcovsky J. 2013. Large deletions in the CBF gene cluster at the Fr-B2 locus are associated with reduced frost tolerance in wheat. *Theor Appl Genet*. 126:2683–2697.
- Pidal B, Yan L, Fu D, Zhang F, Tranquilli G, Dubcovsky J. 2009. The CARG-box located upstream from the transcriptional start of wheat vernalization gene VRN1 Is not necessary for the vernalization response. *J Hered*. 100:355–364.
- Quamme H, Stushnoff C, Weiser CJ. 1972. The relationship of exotherms to cold injury in apple stem tissues. *J Am Soc Hortic Sci*. 97:608–613.
- Rajashekar CB, Lafta A. 1996. Cell-wall changes and cell tension in response to cold acclimation and exogenous abscisic acid in leaves and cell cultures. *Plant Physiol*. 111:605–612.
- Rao KVM, Raghavendra AS, Reddy KJ. 2006. Physiology and molecular biology of stress tolerance in plants. Springer.
- Riechmann J, Heard J, Martin G, Reuber L, Keddie J, Adam L, Pineda O, Ratcliffe O, Samaha R, Creelman R. 2000. Arabidopsis transcription factors: genome-wide comparative analysis among eukaryotes. *Science*. 290:2105–2110.
- Riechmann JL, Meyerowitz EM. 1998. The AP2/EREBP family of plant transcription factors. *Biol Chem*. 379:633–646.
- Rudd JJ, Franklin-Tong VE. 2001. Unravelling response-specificity in Ca²⁺ signalling pathways in plant cells. *New Phytol*. 151:7–33.
- Ríos G, Leida C, Conejero A, Badenes ML. 2014. Epigenetic regulation of bud dormancy events in perennial plants. *Front Plant Sci*. 5:247.
- Sakai A, Larcher W. 1987. Frost survival of plants. Berlin: Springer.
- Sakuma Y, Liu Q, Dubouzet JG, Abe H, Shinozaki K, Yamaguchi-Shinozaki K. 2002. DNA-binding specificity of the ERF/AP2 domain of Arabidopsis DREBs, transcription factors involved in dehydration- and cold-inducible gene expression. *Biochem Biophys Res Commun*. 290:998–1009.
- Sanghera GS, Wani SH, Hussain W, Singh NB. 2011. Engineering cold stress tolerance in crop plants. *Curr Genomics*. 12:30.
- Sattler U, Calsou P, Boiteux S, Salles B. 2000. Detection of oxidative base DNA damage by a new biochemical assay. *Arch Biochem Biophys*. 376:26–33.
- Sayyari M, Babalar M, Kalantari S, Martínez-Romero D, Guillén F, Serrano M, Valero D. 2011. Vapour treatments with methyl salicylate or methyl jasmonate alleviated chilling injury and enhanced antioxidant potential during postharvest storage of pomegranates. *Food Chem*. 124:964–970.
- Seki M, Kamei A, Yamaguchi-Shinozaki K, Shinozaki K. 2003. Molecular responses to drought, salinity and frost: common and different paths for plant protection. *Curr Opin Biotechnol*. 14:194–199.
- Seki M, Narusaka M, Abe H, Kasuga M, Yamaguchi-Shinozaki K, Carninci P, Hayashizaki Y, Shinozaki K. 2001. Monitoring the expression pattern of 1300 Arabidopsis genes under drought and cold stresses by using a full-length cDNA microarray. *Plant Cell*. 13:61–72.
- Seki M, Narusaka M, Ishida J, Nanjo T, Fujita M, Oono Y, Kamiya A, Nakajima M, Enju A, Sakurai T, et al. 2002. Monitoring the expression profiles of 7000 Arabidopsis genes under drought, cold and high-salinity stresses using a full-length cDNA microarray. *Plant J*. 31:279–292.
- Seo PJ, Lee A-K, Xiang F, Park C-M. 2008. Molecular and functional profiling of Arabidopsis pathogenesis-related genes: insights into their roles in salt response of seed germination. *Plant Cell Physiol*. 49:334–344.
- Shinozaki K, Yamaguchi-Shinozaki K. 2000. Molecular responses to dehydration and low temperature: differences and cross-talk between two stress signaling pathways. *Curr Opin Plant Biol*. 3:217–223.
- Soltész A, Smedley M, Vashegyi I, Galiba G, Harwood W, Vágújfalvi A. 2013. Transgenic barley lines prove the involvement of TaCBF14 and TaCBF15 in the cold acclimation process and in frost tolerance. *J Exp Bot*. 64:1849–1862.
- Steponkus PL, Uemura M, Joseph RA, Gilmour SJ, Thomashow MF. 1998. Mode of action of the COR15a gene on the freezing tolerance of *Arabidopsis thaliana*. *Proc Natl Acad Sci USA*. 95:14570–14575.
- Steponkus PL, Uemura M, Webb M. 1993. Membrane destabilization during freeze-induced dehydration. *Curr Top Plant Physiol*. 10:37.
- Stockinger EJ, Skinner JS, Gardner KG, Francia E, Pecchioni N. 2007. Expression levels of barley Cbf genes at the frost resistance-H2 locus are dependent upon alleles at Fr-H1 and Fr-H2. *Plant J*. 51:308–321.
- Stushnoff C, Flower DB, Brule-Babel A. 1984. Breeding and selection for resistance to low-temperature. In: PB Vose, editor. Plant breeding – a contemporary basis. Elsevier; p. 115–136.
- Sun X, Hu C, Tan Q, Gan Q. 2006. Effects of molybdenum on photosynthetic characteristics in winter wheat under low temperature stress. *Zuo wu xue bao*. 32:1418–1422.
- Sun X, Hu C, Tan Q, Liu J, Liu H. 2009. Effects of molybdenum on expression of cold-responsive genes in abscisic acid (ABA)-dependent and ABA-independent pathways in winter wheat under low-temperature stress. *Ann Bot*. 104:345–356.
- Sun X, Zhao T, Gan S, Ren X, Fang L, Karungo SK, Wang Y, Chen L, Li S, Xin H. 2016. Ethylene positively regulates cold tolerance in grapevine by modulating the expression of ethylene response factor 057. *Sci Rep*. 6.
- Takahashi D, Li B, Nakayama T, Kawamura Y, Uemura M. 2013. Plant plasma membrane proteomics for improving cold tolerance. *Front Plant Sci*. 4:90.
- Tao D-L, Öquist G, Wingsle G. 1998. Active oxygen scavengers during cold acclimation of Scots pine seedlings in relation to freezing tolerance. *Cryobiology*. 37:38–45.
- Thomashow MF. 1999. Plant cold acclimation: freezing tolerance genes and regulatory mechanisms. *Annu Rev Plant Physiol Plant Mol Biol*. 50:571–599.
- Thomashow MF. 2001. So what's new in the field of plant cold acclimation? Lots!. *Plant Physiol*. 125:89–93.
- Thomashow MF. 2010. Molecular basis of plant cold acclimation: insights gained from studying the CBF cold response pathway. *Plant Physiol*. 154:571–577.
- Timperio AM, Egidi MG, Zolla L. 2008. Proteomics applied on plant abiotic stresses: role of heat shock proteins (HSP). *J Proteomics*. 71:391–411.
- Ton J, Flors V, Mauch-Mani B. 2009. The multifaceted role of ABA in disease resistance. *Trends Plant Sci*. 14:310–317.
- Tuteja N. 2007. Abscisic acid and abiotic stress signaling. *Plant Signal Behav*. 2:135–138.
- Tyystjärvi E. 2013. Chapter 7, Photoinhibition of Photosystem II. In: Kwang WJ, editor. International review of cell and molecular biology. Academic Press.
- Uemura M, Steponkus PL. 1994. A contrast of the plasma-membrane lipid composition of oat and rye leaves in relation to freezing tolerance. *Plant Physiol*. 104:479–496.
- Uno Y, Furihata T, Abe H, Yoshida R, Shinozaki K, Yamaguchi-Shinozaki K. 2000. Arabidopsis basic leucine zipper transcription factors involved in an abscisic acid-dependent signal transduction pathway under drought and high-salinity conditions. *Proc Natl Acad Sci USA*. 97:11632–11637.
- Vij S, Tyagi AK. 2007. Emerging trends in the functional genomics of the abiotic stress response in crop plants. *Plant Biotechnol J*. 5:361–380.
- Vogel JT, Zarka DG, Van Buskirk HA, Fowler SG, Thomashow MF. 2005. Roles of the CBF2 and ZAT12 transcription factors in configuring the low temperature transcriptome of Arabidopsis. *Plant J*. 41:195–211.
- Vágújfalvi A, Galiba G, Dubcovsky J, Cattivelli L. 2000. Two loci on wheat chromosome 5A regulate the differential cold-dependent expression of the cor14b gene in frost-tolerant and frost-sensitive genotypes. *Mol Gen Genet MGG*. 263:194–200.
- Walters Jr KR, Serianni AS, Voituron Y, Sformo T, Barnes BM, Duman JG. 2011. A thermal hysteresis-producing xylomannan glycolipid antifreeze associated with cold tolerance is found in diverse taxa. *J Comp Physiol B*. 181:631–640.
- Wang Q, Guan Y, Wu Y, Chen H, Chen F, Chu C. 2008. Overexpression of a rice OsDREB1F gene increases salt, drought, and low temperature tolerance in both Arabidopsis and rice. *Plant Mol Biol*. 67:589–602.

- Wani SH, Kumar V, Shriram V, Sah SK. 2016. Phytohormones and their metabolic engineering for abiotic stress tolerance in crop plants. *Crop J.* 4:162–176.
- Wanner LA, Junttila O. 1999. Cold-induced freezing tolerance in *Arabidopsis*. *Plant Physiol.* 120:391–400.
- Wan S-B, Tian L, Tian R-R, Pan Q-H, Zhan J-C, Wen P-F, Chen J-Y, Zhang P, Wang W, Huang W-D. 2009. Involvement of phospholipase D in the low temperature acclimation-induced thermotolerance in grape berry. *Plant Physiol Biochem.* 47:504–510.
- Welling A, Palva ET. 2008. Involvement of CBF transcription factors in winter hardiness in birch. *Plant Physiol.* 147:1199–1211.
- Williams R, Frausto da Silva J. 2002. The involvement of molybdenum in life. *Biochem Biophys Res Commun.* 292:293–299.
- Winfield MO, Lu C, Wilson ID, Coghill JA, Edwards KJ. 2010. Plant responses to cold: transcriptome analysis of wheat. *Plant Biotechnol J.* 8:749–771.
- Wisniewski M, Fuller M. 1999. Ice nucleation and deep supercooling in plants: new insights using infrared thermography. In: *Cold-adapted organisms: ecology, physiology, enzymology and molecular biology*. Berlin: Springer-Verlag; p. 105–118.
- Wisniewski M, Glenn DM, Gusta L, Fuller MP. 2008. Using infrared thermography to study freezing in plants. *HortScience.* 43:1648–1651.
- Wisniewski M, Lindow SE, Ashworth EN. 1997. Observations of ice nucleation and propagation in plants using infrared video thermography. *Plant Physiol.* 113:327–334.
- Wisniewski M, Nassuth A, Teulieres C, Marque C, Rowland J, Cao PB, Brown A. 2014. Genomics of cold hardiness in woody plants. *Crit Rev Plant Sci.* 33:92–124.
- Wisniewski M, Norelli J, Artlip T. 2015. Overexpression of a peach CBF gene in apple: a model for understanding the integration of growth, dormancy, and cold hardiness in woody plants. *Front Plant Sci.* 6:85.
- Xin Z, Browse J. 2000. Cold comfort farm: the acclimation of plants to freezing temperatures. *Plant Cell Environ.* 23:893–902.
- Xiong L, Ishitani M, Lee H, Zhu JK. 2001. The *Arabidopsis* LOS5/ABA3 locus encodes a molybdenum cofactor sulfurase and modulates cold stress- and osmotic stress-responsive gene expression. *Plant Cell Online.* 13:2063–2083.
- Xue-Xuan X, Hong-Bo S, Yuan-Yuan M, Gang X, Jun-Na S, Dong-Gang G, Cheng-Jiang R. 2010. Biotechnological implications from abscisic acid (ABA) roles in cold stress and leaf senescence as an important signal for improving plant sustainable survival under abiotic-stressed conditions. *Crit Rev Biotechnol.* 30:222–230.
- Xu XM, Meier I. 2008. The nuclear pore comes to the fore. *Trends Plant Sci.* 13:20–27.
- Yamaguchi-Shinozaki K, Shinozaki K. 1994. A novel cis-acting element in an *Arabidopsis* gene is involved in responsiveness to drought, low-temperature, or high-salt stress. *Plant Cell.* 6:251–264.
- Yamaguchi-Shinozaki K, Shinozaki K. 2005. Organization of cis-acting regulatory elements in osmotic- and cold-stress-responsive promoters. *Trends Plant Sci.* 10:88–94.
- Yamasaki Y, Randall SK. 2016. Functionality of soybean CBF/DREB1 transcription factors. *Plant Sci.* 246:80–90.
- Yamori W, Noguchi K, Hikosaka K, Terashima I. 2010. Phenotypic plasticity in photosynthetic temperature acclimation among crop species with different cold tolerances. *Plant Physiol.* 152:388–399.
- Yang Z, Chen C, Xiaofen J, Aisheng X, Rihe P, Yihuan H, Quanhong Y, Jianmin C. 2009. Expression of a rice DREB1 gene, OsDREB1D, enhances cold and high-salt tolerance in transgenic *Arabidopsis*. *BMB Rep (Biochem Mol Biol Rep).* 42:486–492.
- Yang W, Liu X-D, Chi X-J, Wu C-A, Li Y-Z, Song L-L, Liu X-M, Wang Y-F, Wang F-W, Zhang C. 2011. Dwarf apple MbDREB1 enhances plant tolerance to low temperature, drought, and salt stress via both ABA-dependent and ABA-independent pathways. *Planta.* 233:219–229.
- Yu M, Hu C, Wang Y. 1999. Influences of seed molybdenum and molybdenum application on nitrate reductase activity, shoot dry matter, and grain yields of winter wheat cultivars. *J Plant Nutr.* 22:1433–1441.
- Zarka DG, Vogel JT, Cook D, Thomashow MF. 2003. Cold induction of *Arabidopsis* CBF genes involves multiple ICE (inducer of CBF expression) promoter elements and a cold-regulatory circuit that is desensitized by low temperature. *Plant Physiol.* 133:910–918.
- Zeller G, Henz SR, Widmer CK, Sachsenberg T, Ratsch G, Weigel D, Laubinger S. 2009. Stress-induced changes in the *Arabidopsis thaliana* transcriptome analyzed using whole-genome tiling arrays. *Plant J.* 58:1068–1082.
- Zhang R, Wang Y, Liu G, Li H. 2010. Cloning and characterization of a pathogenesis-related gene (ThPR10) from *Tamarix hispida*. *Acta Biologica Cracoviensia Series Botanica.* 52:17–25.
- Zhang, J, Xu, Y, Huan, Q, Chong, K. 2009. Deep sequencing of *Brachypodium* small RNAs at the global genome level identifies microRNAs involved in cold stress response. *BMC Genomics.* 10, 449.
- Zhao ML, Wang JN, Shan W, Fan JG, Kuang JF, Wu KQ, Li XP, Chen WX, He FY, Chen JY. 2013. Induction of jasmonate signalling regulators MaMYC2s and their physical interactions with MaICE1 in methyl jasmonate-induced chilling tolerance in banana fruit. *Plant Cell Environ.* 36:30–51.
- Zhao C, Zhu J-K. 2016. The broad roles of CBF genes: from development to abiotic stress. *Plant Signal Behav.* 11:e1215794.
- Zhou X, Wang G, Sutoh K, Zhu J-K, Zhang W. 2008. Identification of cold-inducible microRNAs in plants by transcriptome analysis. *Biochim Biophys Acta (BBA) – Gene Regul Mech.* 1779:780–788.
- Zwack PJ, Rashotte AM. 2015. Interactions between cytokinin signalling and abiotic stress responses. *J Exp Bot.* 66:4863–4871.